Pig IgG Heavy and Light Chain Cross-Adsorbed Antibody



Goat Polyclonal Antigen Affinity Purified Catalog No. A100-205A

Lot No. 2	240123
APPLICATIONS	WB, IHC, ICC, ELISA
SPECIES REACTIVI	TY Pig. Minimum reactivity to bovine, chicken, horse, human, mouse, rabbit and rat
AMOUNT	1 ml
CONCENTRATION	0.5 mg/ml
STORAGE/SHELF L	IFE 2 - 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antiserum was cross adsorbed using bovine, chicken, horse, human, mouse, rabbit and rat immunosorbents to remove cross reactive Antibodies. The antibody to pig IgG was isolated by affinity chromatography using antigen coupled to agarose beads.
	Immunoglobulin concentration was determined using Beer's Law where 1mg/mL IgG has an A280 of 1.4.
	By immunoelectrophoresis and ELISA this antibody reacts specifically with pig IgG and with light chains common to other pig immunoglobulins. No antibody was detected against non- immunoglobulin serum proteins. Less than 1% cross reactivity to bovine, chicken, horse, human, mouse, rabbit and rat IgG was detected. This antibody may cross react with IgG from other species.
APPLICATIONS	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.
	Western Blot1:1000 - 1:20,000Immunohistochemistry1:100 - 1:1000Immunocytochemistry1:100 - 1:1000ELISA1:1000 - 1:20,000; for coating plates 1:50 - 1:250
APPLICATION NOT	TES Not all listed applications have been specifically tested by our laboratory.
ADDITIONAL INFO	https://www.fortislife.com/p/A100-205A Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Michael Spencer, PhD Date: January 24, 2024

1