

Rat IgM cross-adsorbed Antibody

Goat Polyclonal Conjugate Biotin
Antigen Affinity Purified
Catalog No. A110-200B
Lot No. A110-200B-6



| | | | | | | | | | |
|------------------------------|---|--------------|----------------------|----------------------|-----------------|---------------------|---------------|-------|----------------------|
| APPLICATIONS | WB, IHC, ICC, ELISA | | | | | | | | |
| SPECIES REACTIVITY | Rat. Minimum reactivity to human and mouse | | | | | | | | |
| ISOTYPE | IgG | | | | | | | | |
| AMOUNT | 1 ml at 0.5 mg/ml | | | | | | | | |
| STORAGE/SHELF LIFE | 2 – 8° C / 1 year from date of receipt | | | | | | | | |
| PHYSICAL STATE | Liquid | | | | | | | | |
| BUFFER | Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide | | | | | | | | |
| ORIGIN | USA | | | | | | | | |
| PRODUCTION PROCEDURES | <p>Antiserum was solid phase adsorbed to ensure class specificity. Antiserum was cross adsorbed using human and mouse immunosorbents to remove cross reactive antibodies. The antibody to rat IgM was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to biotin.</p> <p>Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.</p> <p>By immunoelectrophoresis and ELISA this antibody reacts specifically with rat IgM. Cross reactivity with IgA and IgG is negligible. No antibody was detected against non-immunoglobulin serum proteins. Less than 1% cross reactivity to human and mouse IgM was detected. This antibody may cross react with IgG from other species.</p> | | | | | | | | |
| APPLICATIONS | <p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <table><tr><td>Western Blot</td><td>1:10,000 – 1:200,000</td></tr><tr><td>Immunohistochemistry</td><td>1:250 – 1:2,500</td></tr><tr><td>Immunocytochemistry</td><td>1:100 – 1:500</td></tr><tr><td>ELISA</td><td>1:10,000 – 1:200,000</td></tr></table> | Western Blot | 1:10,000 – 1:200,000 | Immunohistochemistry | 1:250 – 1:2,500 | Immunocytochemistry | 1:100 – 1:500 | ELISA | 1:10,000 – 1:200,000 |
| Western Blot | 1:10,000 – 1:200,000 | | | | | | | | |
| Immunohistochemistry | 1:250 – 1:2,500 | | | | | | | | |
| Immunocytochemistry | 1:100 – 1:500 | | | | | | | | |
| ELISA | 1:10,000 – 1:200,000 | | | | | | | | |
| APPLICATION NOTES | Not all listed applications have been specifically tested by our laboratory. | | | | | | | | |
| ADDITIONAL INFO | <p>http://www.bethyl.com/product/A110-200B</p> <p>Use the link above to view SDS, a current list of citations, and other product specific information.</p> | | | | | | | | |