Rat IgG Heavy and Light Chain Cross-Adsorbed Antibody



Rabbit Polyclonal Antigen Affinity Purified Catalog No. A110-322A

Lot No.	230428	
APPLICATIONS		WB, IHC, ICC, ELISA
SPECIES REACTI	VITY	Rat. Minimum reactivity to bovine, horse, human, mouse and sheep
AMOUNT		1 ml
CONCENTRATIO	ON	0.5 mg/ml
STORAGE/SHEL	F LIFE	2 - 8°C / 1 year from date of receipt
PHYSICAL STAT	E	Liquid
BUFFER		Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide
ISOTYPE		IgG
ORIGIN		USA
PRODUCTION PROCEDURES		Antiserum was cross adsorbed using bovine, horse, human, mouse & sheep immunosorbents to remove cross reactive antibodies. The antibody to rat IgG was isolated by affinity chromatography using antigen coupled to agarose beads.
		Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. By immunoelectrophoresis and ELISA this antibody reacts specifically with rat IgG and with light chains common to other rat immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. Less than 1% cross reactivity to bovine, horse, human, mouse & sheep IgG was detected. This antibody may cross react with IgG from other species.
APPLICATIONS		Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.
		Western Blot1:1,000 - 1:20,000Immunohistochemistry1:200 - 1:2,000Immunocytochemistry1:200 - 1:2,000ELISA1:1,000 - 1:20,000; for coating plates 1:50 - 1:250
APPLICATION N	OTES	Not all listed applications have been specifically tested by our laboratory.
ADDITIONAL IN	FO	https://www.fortislife.com/p/A110-322A Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Michael Spencer, PhD Date: May 2, 2023