Rabbit IgG Heavy and Light Chain Antibody



Goat Polyclonal Conjugate DyLight® 650

Antigen Affinity Purified

Catalog No. A120-101D5

Lot No. 12

APPLICATIONS IHC, ICC, Flow Cyt, IF

SPECIES REACTIVITY Rabbit

AMOUNT 1 ml

CONCENTRATION 0.5 mg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide

FLUOROPHORE/PROTEIN 5.5
ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

The antibody was isolated by affinity chromatography using antigen coupled to agarose

beads and conjugated to DyLight® 650.

Prior to conjugation, immunoglobulin concentration was determined using Beer's Law where

1mg/mL lgG has an A280 of 1.4.

By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG and with light chains common to other rabbit immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from

other species.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry1:50 - 1:500Immunocytochemistry1:50 - 1:500Flow Cytometry1:50 - 1:200Immunofluorescence1:50 - 1:500

APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.

DyLight® 650 is excited at 652 (in PBS) and emits at 672 (in PBS). DyLight® 650 replaces DyLight®

649.

DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.

ADDITIONAL INFO https://www.fortislife.com/p/A120-101D5


Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: November 20, 2023

Phone: 800.338.9579 • Fax: 866.597.6105 • Web: www.fortislife.com Orders: orders@fortislife.com • Support: technical@fortislife.com



Detection of human E-Cadherin (shaded) in HT29 cells by flow cytometry. *Antibody:* Rabbit anti-E-Cadherin recombinant monoclonal antibody [BLR088G] (A700-088) or isotype control (unshaded). *Secondary:* DyLight® 650-conjugated goat anti-rabbit IgG (A120-101D5).