Rabbit IgG Heavy and Light Chain Antibody



Donkey Polyclonal Conjugate DyLight® 594

Antigen Affinity Purified

Catalog No. A120-108D4

Lot No. A120-108D4-10

APPLICATIONS IHC, ICC, F, IF

SPECIES REACTIVITY Rabbit

AMOUNT 1 ml

CONCENTRATION 0.5 mg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide

FLUOROPHORE/PROTEIN 6.5
ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

The antibody was isolated by affinity chromatography using antigen coupled to agarose

beads and conjugated to DyLight® 594.

Antibody concentration was determined by extinction coefficient prior to conjugation:

absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG and with light chains common to other rabbit immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from

other species.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry 1:50 – 1:500
Immunocytochemistry 1:50 – 1:500
Flow Cytometry 1:50 – 1:200
Immunofluorescence 1:50 – 1:500

APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.

DyLight® 594 is excited at 593 (in PBS) and emits at 618 (in PBS).

DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.

ADDITIONAL INFO https://www.fortislife.com/p/A120-108D4


Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: July 1, 2022

Phone: 800.338.9579 • Fax: 866.597.6105 • Web: www.fortislife.com Orders: orders@fortislife.com • Support: technical@fortislife.com



Detection of human PCNA by immunofluorescence. *Sample:* FFPE section of human stomach carcinoma. *Primary Antibody:* Affinity purified rabbit anti-PCNA (Cat. No. IHC-00012 Lot1) used at a dilution of 1:100. *Secondary Antibody:* Red-fluorescent donkey anti-rabbit IgG-heavy and light chain Antibody DyLight® 594 Conjugated (A120-108D4 Lot 4) used at a dilution of 1:100 (5µg/ml). *Counterstain:* DAPI (blue)