Rabbit IgG Heavy and Light Chain Cross-Adsorbed Antibody



Donkey Polyclonal Conjugate DyLight® 550

Antigen Affinity Purified
Catalog No. A120-208D3

Lot No. 15

APPLICATIONS IHC, ICC, Flow Cyt, IF

SPECIES REACTIVITY Rabbit. Minimum reactivity to bovine, chicken, goat, human, mouse, pig and rat

AMOUNT 1 ml

CONCENTRATION 0.5 mg/ml

STORAGE/SHELF LIFE 2 – 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide

FLUOROPHORE/PROTEIN 4.8
ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antiserum was cross adsorbed using bovine, chicken, goat, human, mouse, pig and rat immunosorbents to remove cross reactive antibodies. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DvLight® 550.

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG and with light chains common to other rabbit immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. Less than 1% cross reactivity to bovine, chicken, goat, human, mouse, pig and rat IgG was detected. This antibody may cross react with IgG from other species.

APPLICATIONS

Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry1:50 - 1:500Immunocytochemistry1:50 - 1:500Flow Cytometry1:50 - 1:200Immunofluorescence1:50 - 1:500

APPLICATION NOTES

Not all listed applications have been specifically tested by our laboratory.

DyLight® 550 is excited at 562 (in PBS) and emits at 576 (in PBS). DyLight® 550 replaces DyLight®

549.

DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.

ADDITIONAL INFO


https://www.fortislife.com/p/A120-208D3

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: April 25, 2023



Detection of human PCNA by immunofluorescence. *Samples:* FFPE sections of human stomach carcinoma. *Primary Antibody:* Affinity purified rabbit anti-PCNA (Cat. No. IHC-00012 Lot1) used at a dilution of 1:100. *Secondary Antibody:* Red-fluorescent donkey anti-rabbit IgG-heavy and light chain cross-adsorbed Antibody DyLight® 550 Conjugated. A120-208D3 Lot 6 (left) and A120-208D3 Lot 7 (right) used at a dilution of 1:100 (5 µg/ml). *Counterstain:* DAPI (blue)