Bat IgG Heavy and Light Chain Antibody



	LABORATORIES, INC. LIFE SCIENCE
Goat Polyclonal	Conjugate Biotin
Antigen Affinity	
5	A140-118B
	9
APPLICATIONS	WB, IHC, ICC, ELISA
SPECIES REACTIV	
AMOUNT	1 ml
CONCENTRATION	
STORAGE/SHELF	
PHYSICAL STATE	Liquid
BUFFER	Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	The antibody was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to biotin.
	Antibody concentration was determined by extinction coefficient prior to conjugation: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. Molar enzyme/antibody protein ratio is 4:1.
	By immunoelectrophoresis and ELISA this antibody reacts specifically with Bat IgG and with light chains common to other Bat immunoglobulins. No antibody was detected against non- immunoglobulin serum proteins. Antibody has been shown to react with bat genus species Pteropus vampirus, Desmodus rotundus, Eptesicus fuscus, Tadrida pumila, T. condylura, Hypsignathus monstrosus, Rosettus aegyptiacus, Epomorphus crypturus, Molossus species, and Phylostomus species.
	This antibody may cross react with IgG from other species.
APPLICATIONS	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.
	Western Blot 1:20,000 – 1:400,000
	Immunohistochemistry 1:500 – 1:5,000
	Immunocytochemistry 1:200 – 1:1,000
	ELISA 1:20,000 - 1:400,000
APPLICATION NO	
ADDITIONAL INF	
	Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Michael Spencer, PhD Date: April 4, 2023

> Phone: 800.338.9579 · Fax: 866.597.6105 · Web: www.fortislife.com Orders: orders@fortislife.com · Support: technical@fortislife.com