## Myc Tag Antibody

Chicken Polyclonal

Antigen Affinity Purified

Catalog No. A190-103A GenelD 4609

Lot No. A190-103A-3

**APPLICATIONS** WB, IP, ICC, ELISA

AMOUNT 0.1 ml
CONCENTRATION 1 mg/ml

STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

**PRODUCTION** Chickens were immunized with a synthetic peptide representing amino acid residues 410–419

PROCEDURES (EQKLISEEDL) of human myc conjugated to KLH. Antibody was isolated by affinity

chromatography using the peptide immobilized on solid support.

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1,000 - 1:30,000

Immunoprecipitation 1 – 4 µg/mg lysate

Immunocytochemistry 1:100 - 1:400

ELISA 1:1,000 – 1:30,000; for coating plates 1:100 – 1:500

**APPLICATION NOTES** Not all listed applications have been specifically tested by our laboratory.


Validation by Western Blot was performed using a Western Blot Gel 4-20%.

ADDITIONAL INFO https://www.bethyl.com/product/A190-103A

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: September 17, 2019





## Detection of c-myc-tagged Protein by western blot.

Samples: 200, 100, or 50 ng of E. coli whole cell lysate expressing a multi-tag fusion protein. Antibodies: Affinity purified, chicken anti-Myc Tag antibody A190-103A used for WB at 0.2  $\mu$ g/ml (1:5,000). Detection: Chemiluminescence with an exposure time of 10 seconds.