Myc Tag Antibody



Goat Polyclonal

Antigen Affinity Purified

Catalog No. A190-104A Uniprot ID P01106 Lot No. A190-104A-221227 GeneID 4609

APPLICATIONS WB, IP, ICC, ELISA

AMOUNT 0.1 ml
CONCENTRATION 1 mg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Goats were immunized with a synthetic peptide representing amino acid residues 410–419

PROCEDURES (EQKLISEEDL) of human myc conjugated to KLH. Antibody was isolated by affinity

chromatography using the peptide immobilized on solid support.

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of

1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1,000 – 1:30,000

Immunoprecipitation 1 – 4 µg/mg lysate

Immunocytochemistry 1:100 - 1:400

ELISA 1:1,000 – 1:30,000; for coating plates 1:100 – 1:500

APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.

Validation by Western Blot was performed using a Western Blot Gel 4-20%.


ADDITIONAL INFO https://www.fortislife.com/p/A190-104A

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: February 22, 2023



Detection of c-myc-tagged Protein by western blot.

Samples: 200, 100, or 50 ng of E. coli whole cell lysate expressing a multi-tag fusion protein. *Antibodies:* Affinity purified, goat anti-Myc Tag antibody A190-104A used for WB at 0.04 μ g/ml (1:25,000). *Detection:* Chemiluminescence with an exposure time of 30 seconds.