Myc Tag Antibody

Goat Polyclonal | Conjugate HRP
Antigen Affinity Purified
Catalog No. A190-104P | Uniprot ID P01106
Lot No. 27 | GeneID 4609

APPLICATIONS	WB, IHC, ICC, ELISA
AMOUNT	0.1 ml
CONCENTRATION	1 mg/ml
STORAGE/SHELF LIFE	2 - 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.05% Pro-Clean 400
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Goats were immunized with a synthetic peptide representing amino acid residues 410-419 (EQKLISEEDL) of human myc conjugated to KLH. Antibody was isolated by affinity chromatography using the peptide immobilized on solid support and conjugated to horseradish peroxidase (HRP). Antibody concentration was determined by extinction coefficient prior to conjugation: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. Molar enzyme/antibody protein ratio is 4:1.
APPLICATIONS	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot	1:1,000 - 1:30,000
Immunohistochemistry	Not applicable
Immunocytochemistry	1:200 - 1:500
ELISA	1:10,000 - 1:100,000

APPLICATION NOTES	Not all listed applications have been specifically tested by our laboratory. Validation by Western Blot was performed using a Western Blot Gel 4-20%.
ADDITIONAL INFO	https://www.fortislife.com/p/A190-104P Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: April 19, 2023



Detection of c-myc-tagged Protein by western blot.
Samples: 200, 100, or 50 ng of E. coli whole cell lysate expressing a multi-tag fusion protein. *Antibodies:* Affinity-purified, HRP-conjugated, goat anti-Myc Tag antibody A190-104P used for WB at 0.04 µg/ml (1:25,000). *Detection:* Chemiluminescence with an exposure time of 10 seconds.