RFC5 Antibody

Goat Polyclonal

Antigen Affinity Purified Protein ID NP_031396.1

Catalog No. A300-146A GeneID 5985

Lot No. A300-146A-2

APPLICATIONS WB

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to RFC5 immobilized on solid support.

The epitope recognized by A300–146A maps to a region between residues 1 and 50 of human

Replication Factor C subunit 5 using the numbering given in entry NP_031396.1 (GeneID 5985).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:500 - 1:2,500

APPLICATION NOTES Western blot of lysates performed using standard western blot reagents and 4-20% SDS-PAGE.

ADDITIONAL INFO https://www.bethyl.com/product/A300-146A


Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019



RFC5 Antibody A300-146A



Detection of human RFC5 by western blot. Samples: Whole cell lysate (50 μ g) from HeLa, HEK293T, and Jurkat cells prepared using NETN lysis buffer. Antibodies: Affinity purified goat anti-RFC5 antibody A300-146A (lot A300-146A-2) used for WB at 1 μ g/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.