ARHGEF6 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_004831.1

Catalog No. A302-558A GeneID 9459

Lot No. A302-558A-1

APPLICATIONS WB, IP
SPECIES REACTIVITY Human
AMOUNT 100 μI

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to ARHGEF6 immobilized on solid

PROCEDURES support.

The epitope recognized by A302-558A maps to a region between residue 50 and 100 of human

Rac/Cdc42 guanine nucleotide exchange factor 6 using the numbering given in entry

NP_004831.1 (GeneID 9459).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000

Immunoprecipitation 5 – 15 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.


ADDITIONAL INFO https://www.bethyl.com/product/A302-558A

Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019





Detection of human ARHGEF6 by western blot and immunoprecipitation. Samples: Whole cell lysate (5, 15 and 50 μg for WB; 1 mg for IP, 20% of IP loaded) from Jurkat cells. Antibodies: Affinity purified rabbit anti–ARHGEF6 antibody A302–558A used for WB at 0.1 μg/ml (A) and 1 μg/ml (B) and used for IP at 10 μg/mg lysate. ARHGEF6 was also immunoprecipitated by rabbit anti–ARHGEF6 antibody A302–559A, which recognizes a downstream epitope. Detection: Chemiluminescence with exposure times of 3 minutes (A) and 10 seconds (B).