FYCO1 Antibody



Antigen Affinity Purified Protein ID NP_078789.2

Catalog No. A302-795A-T GeneID 79443

Lot No. A302-795A-T-1

APPLICATIONS WB, IP

SPECIES REACTIVITY Human

AMOUNT 10 μl

CONCENTRATION 200 μg/ml

STORAGE/SHELF LIFE 2 – 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to FYCO1 immobilized on solid

PROCEDURES support.

The epitope recognized by A302–795A–T maps to a region between residue 325 and 375 of

human FYVE and Coiled-Coil Domain Containing 1 using the numbering given in entry

NP_078789.2 (GeneID 79443).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2.000 - 1:10.000

Immunoprecipitation $5 - 15 \mu g/mg$ lysate

ADDITIONAL INFO https://www.bethyl.com/product/A302-795A-T

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: June 6, 2022

Phone: 800.338.9579 • Fax: 866.597.6105 • Web: www.bethyl.com Orders: orders@fortislife.com • Support: technical@fortislife.com FYCO1 Antibody A302-795A-T



Detection of human FYCO1 by western blot and immunoprecipitation. *Samples:* Whole cell lysate from HeLa (5, 15 and 50 μg for WB; 1 mg for IP, 20% of IP loaded) and HEK293T (T; 50 μg) cells. *Antibodies:* Affinity purified rabbit anti–FYCO1 antibody A302–795A used for WB at 0.04 μg/ml (A) and 0.4 μg/ml (B) and used for IP at 10 μg/mg lysate. FYCO1 was also immunoprecipitated by rabbit anti–FYCO1 antibody A302–796A, which recognizes a downstream epitope. *Detection:* Chemiluminescence with exposure times of 3 minutes (A) and 30 seconds (B).