

E2F8 Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Catalog No. A303-039A


Lot No. A303-039A-1

Protein ID NP_078956.2

Gene ID 79733



APPLICATIONS	WB, IP				
REACTIVITY TESTED	Human, Mouse				
PRESUMED REACTIVITY	Based on 100% sequence identity, this antibody is predicted to react with Rat, Dog, Guinea pig_10141, Panda, Orangutan, Rhesus Monkey, Gorilla, Chimpanzee, Tasmanian devil, White-tufted-ear marmoset, Crab-eating macaque, Chinese hamster, Naked mole rat and Northern white-cheeked gibbon.				
ISOTYPE	IgG				
AMOUNT	0.1 ml at 1 mg/ml				
STORAGE/SHELF LIFE	2 - 8° C / 1 year from date of receipt				
PHYSICAL STATE	Liquid				
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide				
ORIGIN	USA				
PRODUCTION PROCEDURES	<p>Antibody was affinity purified using an epitope specific to E2F8 immobilized on solid support.</p> <p>The epitope recognized by A303-039A maps to a region between residue 817 and 867 of human E2F Transcription Factor 8 using the numbering given in entry NP_078956.2 (GeneID 79733).</p> <p>Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.</p>				
APPLICATIONS	<p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <table><tr><td>Western Blot</td><td>1:2,000 to 1:10,000</td></tr><tr><td>Immunoprecipitation</td><td>2 to 5 µg/mg lysate</td></tr></table>	Western Blot	1:2,000 to 1:10,000	Immunoprecipitation	2 to 5 µg/mg lysate
Western Blot	1:2,000 to 1:10,000				
Immunoprecipitation	2 to 5 µg/mg lysate				
APPLICATION NOTES	Validation by IP/Western Blot was performed using a 4-8% SDS-PAGE gel and ReliaBLOT® Reagents (Cat. No. WB120).				
ADDITIONAL INFO	<p>http://www.bethyl.com/product/A303-039A</p> <p>Use the link above to view SDS, a current list of citations, and other product specific information.</p>				



Detection of Human and Mouse E2F8 by Western Blot (h&m) and Immunoprecipitation (h). *Samples:* Whole cell lysate from HeLa (H; 50 μ g for WB; 1 mg for IP, 20% of IP loaded), 293T (5, 15, and 50 μ g for WB) and mouse NIH3T3 (M; 50 μ g) cells. *Antibodies:* Affinity purified rabbit anti-E2F8 antibody A303-039A used for WB at 0.1 μ g/ml (A) and 1 μ g/ml (B) and used for IP at 6 μ g/mg lysate. E2F8 was also immunoprecipitated by rabbit anti-E2F8 antibody BL11013, which recognizes an upstream epitope. *Detection:* Chemiluminescence with exposure times of 3 minutes (A) and 10 seconds (B).