## **AS160 Antibody**



Antigen Affinity Purified Protein ID NP\_055647.2

Catalog No. A303-727A-T GeneID 9882

Lot No. A303-727A-T-1

APPLICATIONS WB, IP
SPECIES REACTIVITY Human
AMOUNT 10 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

**PRODUCTION** Antibody was affinity purified using an epitope specific to AS160 immobilized on solid

**PROCEDURES** support.

The epitope recognized by A303-727A-T maps to a region between residue 1 and 50 of human Akt Substrate of 160 kDa using the numbering given in entry NP\_055647.2 (GeneID

9882).

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2.000 - 1:10.000

Immunoprecipitation  $2 - 10 \mu g/mg$  lysate

ADDITIONAL INFO https://www.bethyl.com/product/A303-727A-T

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol\_IP\_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: June 6, 2022

Phone: 800.338.9579 • Fax: 866.597.6105 • Web: www.bethyl.com Orders: orders@fortislife.com • Support: technical@fortislife.com AS160 Antibody A303-727A-T



## Detection of human AS160 by western blot and immunoprecipitation. Samples: Whole cell lysate from HeLa (15 and 50 $\mu$ g for WB; 1 mg for IP, 20% of IP loaded), HEK293T (T; 50 $\mu$ g) and Jurkat (J; 50 $\mu$ g) cells. Antibodies: Affinity purified rabbit anti-AS160 antibody A303-727A used for WB at 0.1 $\mu$ g/ml (A) and 1 $\mu$ g/ml (B) and used for IP at 6 $\mu$ g/mg lysate. AS160 was also immunoprecipitated by rabbit anti-AS160 antibodies A303-728A and A303-729A, which recognize downstream epitopes. Detection: Chemiluminescence with exposure times of 3 minutes (A) and 30 seconds (B).