

CHAMP1 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_115812.1

Catalog No. A304-216A-T GenelD 283489

Lot No. A304-216A-T-1

APPLICATIONS	WB, IP
SPECIES REACTIVITY	Human, Mouse
AMOUNT	10 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to CHAMP1 immobilized on solid support. The epitope recognized by A304-216A-T maps to a region between residue 1 to 50 of human Chromosome Alignment Maintaining Phosphoprotein 1 using the numbering given in entry NP_115812.1 (GenelD 283489).


APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 – 1:10,000

Immunoprecipitation 2 – 10 µg/mg lysate

ADDITIONAL INFO <https://www.bethyl.com/product/A304-216A-T>
Use the link above to view SDS, a current list of citations, and other product specific information.
IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: June 6, 2022




Detection of human and mouse CHAMP1 by western blot.

Samples: Whole cell lysate (50 µg) from HeLa, HEK293T, Jurkat, mouse TCMK-1, and mouse NIH 3T3 cells.

Antibodies: Affinity purified rabbit anti-CHAMP1 antibody A304-216A (lot A304-216A-1) used for WB at 0.1 µg/ml.

Detection: Chemiluminescence with an exposure time of 3 minutes.



Detection of human CHAMP1 by western blot of immunoprecipitates.

Samples: Whole cell lysate (1 mg for IP; 20% of IP loaded) from 293T cells. *Antibodies:* Affinity purified rabbit anti-CHAMP1 antibody A304-216A (lot A304-216A-1) used for IP at 6 µg/mg lysate. CHAMP1 was also immunoprecipitated by rabbit anti-CHAMP1 antibody A304-217A. For blotting immunoprecipitated CHAMP1, A304-216A was used at 1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.