

# MYCL1 /L-Myc Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID P12524

Catalog No. A304-795A-T GenelD 4610

Lot No. A304-795A-T-1

---

<b>APPLICATIONS</b>	IP
<b>SPECIES REACTIVITY</b>	Human
<b>AMOUNT</b>	10 µl
<b>CONCENTRATION</b>	1000 µg/ml
<b>STORAGE/SHELF LIFE</b>	2 – 8°C / 1 year from date of receipt
<b>PHYSICAL STATE</b>	Liquid
<b>BUFFER</b>	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
<b>ISOTYPE</b>	IgG
<b>ORIGIN</b>	USA
<b>PRODUCTION PROCEDURES</b>	Antibody was affinity purified using an epitope specific to MYCL1 /L-Myc immobilized on solid support.

The epitope recognized by A304-795A-T maps to a region between residue 125 to 175 of human V-Myc Avian Myelocytomatosis Viral Oncogene Lung Carcinoma Derived Homolog using the numbering given in entry P12524 (GenelD 4610).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended. A304-795A has not performed satisfactorily when used for WB of MYCL1 /L-Myc in crude preparations (e.g. whole cell lysate). This antibody can be used for WB of enriched (e.g. immunoprecipitated) sources of MYCL1 /L-Myc.


Immunoprecipitation 2 – 10 µg/mg lysate

**ADDITIONAL INFO** <https://www.bethyl.com/product/A304-795A-T>

Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: [https://www.bethyl.com/content/protocol\\_IP\\_WB](https://www.bethyl.com/content/protocol_IP_WB)

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.  
Michael Spencer, PhD Date: June 6, 2022



**Detection of human MYCL1/L-Myc by western blot of immunoprecipitates.** *Samples:* Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from 293T cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-MYCL1/L-Myc antibody A304-795A (lot A304-795A-1) used for IP at 6 µg per reaction. MYCL1/L-Myc was also immunoprecipitated by rabbit anti-MYCL1/L-Myc antibody A304-796A. For blotting immunoprecipitated MYCL1/L-Myc, A304-795A was used at 0.4 µg/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.