

YARS/Tyrosyl-tRNA Synthetase/TYRRS Antibody



Rabbit Polyclonal				
Antigen Affinity Purified		Protein ID	P54577.4	
Catalog No. A305-0		064A-T	GenelD	8565
Lot No.	A305-	064A-T-1		
APPLICATIONS		WB, IP		
SPECIES REACTIVITY		Human, Mouse		
PRESUMED REACTIVITY		Based on 100% sequence identity, this antibody is predicted to react with Rat and Orangutan		
AMOUNT		10 μΙ		
CONCENTRATION		1000 μg/ml		
STORAGE/SHELF LIFE		2 - 8°C / 1 year from date of receipt		
PHYSICAL STATE		Liquid		
BUFFER		Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide		
ISOTYPE		IgG		
ORIGIN		USA		
PRODUCTION PROCEDURES		Antibody was affinity purified using an epitope specific to YARS/Tyrosyl-tRNA Synthetase/TYRRS immobilized on solid support.		
APPLICATIONS		The epitope recognized by A305-064A-T maps to a region between residue 1 to 50 of human Tyrosyl-tRNA synthetase, cytoplasmic using the numbering given in entry P54577.4 (GeneID 8565).		
		Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.		
		Western Blot	1:2	2,000 - 1:10,000
ADDITIONAL INFO		Immunoprecip	itation 2 -	10 µg/mg lysate
		https://www.bethyl.com/product/A305-064A-T Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB		

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Michael Spencer, PhD Date: June 6, 2022





Detection of human and mouse YARS by western blot. Samples: Whole cell lysate (50 μ g) from HeLa, HEK293T, Jurkat, mouse TCMK-1, and mouse NIH 3T3 cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-YARS antibody A305-064A (lot A305-064A-1) used for WB at 0.1 μ g/ml. Detection: Chemiluminescence with an exposure time of 30 seconds. Detection of human YARS by western blot of immunoprecipitates. *Samples:* Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from 293T cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti–YARS antibody A305–064A (lot A305–064A–1) used for IP at 6 µg per reaction. YARS was also immunoprecipitated by rabbit anti–YARS antibody BL19937. For blotting immunoprecipitated YARS, A305–064A was used at 0.4 µg/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.