## DIEXF/C1 or f107 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID Q68CQ4.2

Catalog No. A305-123A GeneID 27042

Lot No. A305-123A-1

APPLICATIONS IP

**SPECIES REACTIVITY** Human 100 μl

CONCENTRATION 1000 μg/ml

**STORAGE/SHELF LIFE** 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

**PRODUCTION** Antibody was affinity purified using an epitope specific to DIEXF/C1orf107 immobilized on solid

**PROCEDURES** support.

The epitope recognized by A305-123A maps to a region between residue 150 to 200 of human

Digestive organ expansion factor homolog using the numbering given in entry Q68CQ4.2

(GeneID 27042).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended. Use rabbit anti–DIEXF antibody A305–122A.

Immunoprecipitation 2 – 10 μg/mg lysate

**APPLICATION NOTES** Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100–020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).


ADDITIONAL INFO https://www.bethyl.com/product/A305-123A

Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol IP WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019





Detection of human DIEXF by western blot of immunoprecipitates. Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from 293T cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-DIEXF antibody A305-123A (lot A305-123A-1) used for IP at 6 μg per reaction. DIEXF was also immunoprecipitated by rabbit anti-DIEXF antibody A305-122A. For blotting immunoprecipitated DIEXF, A305-123A was used at 0.4 μg/ml. Detection: Chemiluminescence with an exposure time of 10 seconds.