## **CD3D Antibody**

Rabbit Polyclonal

Antigen Affinity Purified Protein ID P04234.1

Catalog No. A305-169A-M GenelD 915

Lot No. A305-169A-M-1

APPLICATIONS WB, IP
SPECIES REACTIVITY Human

**AMOUNT** 100 μl (10 blots)

CONCENTRATION 100 μg/ml

**STORAGE/SHELF LIFE** 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to CD3D immobilized on solid support.

The epitope recognized by A305-169A-M maps to a region between residue 121 to 171 of human T-cell surface glycoprotein CD3 delta chain using the numbering given in entry P04234.1

(GeneID 915).

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1000

Immunoprecipitation The antibody contained within A305-169A-M has been qualified for

use in immunoprecipitation; however, we recommend using the alternative formulation of this antibody found as product A305-169A.

**APPLICATION NOTES** Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100–020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-12% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-12% SDS-PAGE.

ADDITIONAL INFO https://www.bethyl.com/product/A305-169A-M

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol\_IP\_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019



CD3D Antibody A305-169A-M



Detection of human CD3D by western blot. Samples: Whole cell lysate (50 μg) from HeLa, HEK293T, and Jurkat cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-CD3D antibody A305–169A–M (lot A305–169A–M–1) used for WB at 1:1000. Detection: Chemiluminescence with an exposure time of 30 seconds.