CD3D Antibody





Rabbit Polyclonal

Antigen Affinity Purified Protein ID P04234.1

Catalog No. A305-169A-T GeneID 915

Lot No. A305-169A-T-1

APPLICATIONS WB, IP
SPECIES REACTIVITY Human
AMOUNT 10 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to CD3D immobilized on solid

PROCEDURES support.

The epitope recognized by A305-169A-T maps to a region between residue 121 to 171 of human T-cell surface glycoprotein CD3 delta chain using the numbering given in entry

P04234.1 (GeneID 915).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000

Immunoprecipitation $2 - 10 \mu g/mg$ lysate

ADDITIONAL INFO https://www.bethyl.com/product/A305-169A-T

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: June 6, 2022

CD3D Antibody A305-169A-T





Detection of human CD3D by western blot. Samples: Whole cell lysate (50 μg) from HeLa, HEK293T, and Jurkat cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-CD3D antibody A305–169A (lot A305–169A–1) used for WB at 0.1 μg/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.

Detection of human CD3D by western blot of immunoprecipitates. Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from Jurkat cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-CD3D antibody A305–169A (lot A305–169A–1) used for IP at 6 μg per reaction. CD3D was also immunoprecipitated by rabbit anti-CD3D antibody BL20487. For blotting immunoprecipitated CD3D, A305–169A was used at 1 μg/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.