

ARHGAP4 Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID P98171.2

Catalog No. A305-232A

GeneID 393

Lot No. A305-232A-1



APPLICATIONS WB, IP

SPECIES REACTIVITY Human

AMOUNT 100 µl

CONCENTRATION 1000 µg/ml

STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG

ORIGIN USA

PRODUCTION PROCEDURES Antibody was affinity purified using an epitope specific to ARHGAP4 immobilized on solid support.

The epitope recognized by A305-232A maps to a region between residue 700 to 750 of human Rho GTPase-activating protein 4 using the numbering given in entry P98171.2 (GeneID 393).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000

Immunoprecipitation 2 - 10 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).


Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.

ADDITIONAL INFO <https://www.bethyl.com/product/A305-232A>


Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019



Detection of human ARHGAP4 by western blot. *Samples:* Whole cell lysate (5 and 15 μ g) from Jurkat cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-ARHGAP4 antibody A305-232A (lot A305-232A-1) used for WB at 0.1 μ g/ml. *Detection:* Chemiluminescence with an exposure time of 3 minutes.



Detection of human ARHGAP4 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from Jurkat cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-ARHGAP4 antibody A305-232A (lot A305-232A-1) used for IP at 6 μ g per reaction. ARHGAP4 was also immunoprecipitated by rabbit anti-ARHGAP4 antibody A305-231A. For blotting immunoprecipitated ARHGAP4, A305-232A was used at 0.4 μ g/ml. *Detection:* Chemiluminescence with an exposure time of 3 minutes.