

# AGTPBP1 Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID Q9UPW5.3

Catalog No. A305-295A-T

GeneID 23287

Lot No. A305-295A-T-1

---

<b>APPLICATIONS</b>	WB, IP
<b>SPECIES REACTIVITY</b>	Human
<b>AMOUNT</b>	10 µl
<b>CONCENTRATION</b>	1000 µg/ml
<b>STORAGE/SHELF LIFE</b>	2 – 8°C / 1 year from date of receipt
<b>PHYSICAL STATE</b>	Liquid
<b>BUFFER</b>	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
<b>ISOTYPE</b>	IgG
<b>ORIGIN</b>	USA
<b>PRODUCTION PROCEDURES</b>	Antibody was affinity purified using an epitope specific to AGTPBP1 immobilized on solid support.

The epitope recognized by A305-295A-T maps to a region between residue 1176 to 1226 of human Cytosolic carboxypeptidase 1 using the numbering given in entry Q9UPW5.3 (GeneID 23287).

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.


Western Blot 1:500 – 1:2,500

Immunoprecipitation 2 – 10 µg/mg lysate


**ADDITIONAL INFO** <https://www.bethyl.com/product/A305-295A-T>

Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol: [https://www.bethyl.com/content/protocol\\_IP\\_WB](https://www.bethyl.com/content/protocol_IP_WB)

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.  
Michael Spencer, PhD Date: June 6, 2022



**Detection of human AGTPBP1 by western blot.** *Samples:* Whole cell lysate (50 µg) from HeLa, 293T, and Jurkat cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-AGTPBP1 antibody A305-295A (lot A305-295A-1) used for WB at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.



**Detection of human AGTPBP1 by western blot of immunoprecipitates.** *Samples:* Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from 293T cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-AGTPBP1 antibody A305-295A (lot A305-295A-1) used for IP at 6 µg per reaction. AGTPBP1 was also immunoprecipitated by rabbit anti-AGTPBP1 antibody BL20934. For blotting immunoprecipitated AGTPBP1, A305-295A was used at 1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.