

DHCR7 Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID Q9UBM7.1

Catalog No. A305-321A

GeneID 1717

Lot No. A305-321A-1



APPLICATIONS	IP
SPECIES REACTIVITY	Human
AMOUNT	100 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 - 8° C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to DHCR7 immobilized on solid support.

The epitope recognized by A305-321A maps to a region between residue 1 to 50 of human 7-dehydrocholesterol reductase using the numbering given in entry Q9UBM7.1 (GeneID 1717).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended. A305-321A has not performed satisfactorily when used for WB of DHCR7 in crude preparations (e.g. whole cell lysate). This antibody can be used for WB of enriched (e.g. immunoprecipitated) sources of DHCR7.

Immunoprecipitation 2 - 10 µg/mg lysate


APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-20% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).

ADDITIONAL INFO <https://www.bethyl.com/product/A305-321A>

Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019



Detection of human DHCR7 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from 293T cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-DHCR7 antibody A305-321A (lot A305-321A-1) used for IP at 6 μ g per reaction. DHCR7 was also immunoprecipitated by rabbit anti-DHCR7 antibody BL21078. For blotting immunoprecipitated DHCR7, A305-321A was used at 1 μ g/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.