

ASPH Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID Q12797.3

Catalog No. A305-342A-T GeneID 444

Lot No. A305-342A-T-1

APPLICATIONS	WB, IP
SPECIES REACTIVITY	Human
AMOUNT	10 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to ASPH immobilized on solid support.

The epitope recognized by A305-342A-T maps to a region between residue 300 to 350 of human Aspartyl/asparaginyl beta-hydroxylase using the numbering given in entry Q12797.3 (GeneID 444).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.


Western Blot 1:2,000 – 1:10,000

Immunoprecipitation 2 – 10 µg/mg lysate


ADDITIONAL INFO <https://www.bethyl.com/product/A305-342A-T>

Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: June 6, 2022



Detection of human ASPH by western blot. *Samples:* Whole cell lysate (50 µg) from HeLa, HEK293T, and Jurkat cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-ASPH antibody A305-342A (lot A305-342A-1) used for WB at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.



Detection of human ASPH by western blot of immunoprecipitates. *Samples:* Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-ASPH antibody A305-342A (lot A305-342A-1) used for IP at 6 µg per reaction. ASPH was also immunoprecipitated by rabbit anti-ASPH antibody A305-343A. For blotting immunoprecipitated ASPH, A305-343A was used at 0.4 µg/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.