

NDUFB6 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID O95139.3

Catalog No. A305-436A-T GenelD 4712

Lot No. A305-436A-T-1

APPLICATIONS	WB, IP
SPECIES REACTIVITY	Human, Mouse
PRESUMED REACTIVITY	Based on 100% sequence identity, this antibody is predicted to react with Bovine, Orangutan, Gorilla and Chimpanzee
AMOUNT	10 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 - 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to NDUFB6 immobilized on solid support.

The epitope recognized by A305-436A-T maps to a region between residue 78 to 128 of human NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 6 using the numbering given in entry O95139.3 (GeneID 4712).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.


Western Blot 1:500 - 1:2,500

Immunoprecipitation 2 - 10 µg/mg lysate

ADDITIONAL INFO <https://www.bethyl.com/product/A305-436A-T>


Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: June 6, 2022



Detection of human and mouse NDUFB6 by western blot.

Samples: Whole cell lysate (15 µg) from HeLa, HEK293T, Jurkat, mouse TCMK-1, and mouse NIH 3T3 cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-NDUFB6 antibody A305-436A (lot A305-436A-1) used for WB at 0.04 µg/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.



Detection of human NDUFB6 by western blot of immunoprecipitates.

Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from 293T cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-NDUFB6 antibody A305-436A (lot A305-436A-1) used for IP at 6 µg per reaction. NDUFB6 was also immunoprecipitated by rabbit anti-NDUFB6 antibody A305-435A. For blotting immunoprecipitated NDUFB6, A305-436A was used at 1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.