

HIGD1A Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID Q9Y241.1

Catalog No. A305-549A-T

GeneID 25994

Lot No. A305-549A-T-1

APPLICATIONS	WB, IP
SPECIES REACTIVITY	Human
PRESUMED REACTIVITY	Based on 100% sequence identity, this antibody is predicted to react with Orangutan
AMOUNT	10 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to HIGD1A immobilized on solid support.

The epitope recognized by A305-549A-T maps to a region between residue 1 to 50 of human HIG1 domain family member 1A using the numbering given in entry Q9Y241.1 (GeneID 25994).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.


Western Blot 1:2,000 – 1:10,000

Immunoprecipitation 2 – 10 µg/mg lysate


ADDITIONAL INFO <https://www.bethyl.com/product/A305-549A-T>

Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: June 6, 2022



Detection of human HIGD1A by western blot. *Samples:* Whole cell lysate (50 μ g) from HeLa, HEK293T, and Jurkat cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-HIGD1A antibody A305-549A (lot A305-549A-1) used for WB at 0.1 μ g/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.



Detection of human HIGD1A by western blot of immunoprecipitates. *Samples:* Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from 293T cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-HIGD1A antibody A305-549A (lot A305-549A-1) used for IP at 6 μ g per reaction. HIGD1A was also immunoprecipitated by rabbit anti-HIGD1A antibody BL22025. For blotting immunoprecipitated HIGD1A, A305-549A was used at 1 μ g/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.