

Goat IgG Heavy and Light Chain Antibody

Rabbit Polyclonal
Antigen Affinity Purified
Catalog No. A50-100D2
Lot No. 4

APPLICATIONS	WB, IHC, ICC, Flow Cyt, IF
SPECIES REACTIVITY	Goat
AMOUNT	1 ml
CONCENTRATION	0.5 mg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide
FLUOROPHORE/PROTEIN	5.7
ISOTYPE	IgG
ORIGIN	USA

PRODUCTION PROCEDURES The antibody was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight® 488.

Prior to conjugation, immunoglobulin concentration was determined using Beer's Law where 1 mg/mL IgG has an A280 of 1.4.

By immunoelectrophoresis and ELISA this antibody reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins.

This antibody may cross react with IgG from other species.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1000 – 1:20,000. 5% non-fat dry milk in PBST or TBST is recommended for blocking and incubation of antibodies. BSA is not recommended.

Immunohistochemistry 1:50 – 1:500

Immunocytochemistry 1:50 – 1:500

Flow Cytometry 1:50 – 1:200

Immunofluorescence 1:50 – 1:500

APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.


DyLight® 488 is excited at 493 (in PBS) and emits at 518 (in PBS).

DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.


ADDITIONAL INFO <https://www.fortislife.com/p/A50-100D2>

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: January 19, 2024



Detection of GAPDH and hnRNP in HeLa Whole Cell Lysate. *Primary Antibodies:* cocktail of goat anti-GAPDH A303-878A (A303-878A-1) and mouse anti-hnRNP A500-011A (A500-011A-1) at 1 µg/ml each. *Secondary Antibodies:* cocktail of Dylight® 488-conjugated rabbit anti-goat A50-100D2 (A50-100D2-2) (blue) and Dylight® 680-conjugated rabbit anti-mouse A90-317D6 (A90-317D6-1) (red) at 0.5 µg/ml each. *Acquisition:* Syngene G:Box, 52 seconds (blue) and 77 seconds (red).



Detection of human CD137 (shaded) in HDLM-2 cells by flow cytometry. *Antibody:* Goat anti-CD137 antibody or isotype control (unshaded). *Secondary:* DyLight® 488-conjugated rabbit anti-goat IgG (A50-100D2).