

Goat IgG Heavy and Light Chain Cross-Adsorbed Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Catalog No. A50-200A

Lot No. 230522

APPLICATIONS	WB, IHC, ICC, ELISA
SPECIES REACTIVITY	Goat. Minimum reactivity to chicken, horse, human, mouse, pig and rat
AMOUNT	1 ml
CONCENTRATION	0.5 mg/ml
STORAGE/SHELF LIFE	2 – 8°C / 2 years from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antiserum was cross adsorbed using chicken, horse, human, mouse, pig and rat immunosorbents to remove cross reactive Antibodies. The antibody to goat IgG was isolated by affinity chromatography using antigen coupled to agarose beads.

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

By immunoelectrophoresis and ELISA this antibody reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. Less than 1% cross reactivity to chicken, horse, human, mouse, pig and rat IgG was detected.

This antibody may cross react with IgG from other species.

APPLICATIONS	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.
	Western Blot 1:1,000 – 1:20,000
	Immunohistochemistry 1:100 – 1:1,000
	Immunocytochemistry 1:100 – 1:1,000
	ELISA 1:1,000 – 1:20,000; for coating plates 1:50 – 1:250

APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.

ADDITIONAL INFO <https://www.fortislife.com/p/A50-200A>
Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: May 23, 2023