

Goat IgG Heavy and Light Chain Cross-Adsorbed Antibody

Donkey Polyclonal Conjugate DyLight® 488

Antigen Affinity Purified

Catalog No. A50-201D2

Lot No. 15

APPLICATIONS	WB, IHC, ICC, Flow Cyt, IF
SPECIES REACTIVITY	Goat. Minimum reactivity to chicken, human, mouse, pig, rabbit and rat
AMOUNT	1 ml
CONCENTRATION	0.5 mg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide
FLUOROPHORE/PROTEIN	6.3
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antiserum was cross adsorbed using chicken, human, mouse, pig, rabbit and rat immunosorbents to remove cross reactive antibodies. The antibody to goat IgG was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight™ 488.

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

By immunoelectrophoresis and ELISA this antibody reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. Less than 1% cross reactivity to chicken, human, mouse, pig, rabbit and rat IgG was detected.

This antibody may cross react with IgG from other species.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot	1:1,000 – 1:20,000. 5% non-fat dry milk in PBST or TBST is recommended for blocking and incubation of antibodies. BSA is not recommended.
Immunohistochemistry	1:50 – 1:500
Immunocytochemistry	1:50 – 1:500
Flow Cytometry	1:50 – 1:200
Immunofluorescence	1:50 – 1:500

APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.

DyLight® 488 is excited at 493 (in PBS) and emits at 518 (in PBS).


DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.

ADDITIONAL INFO <https://www.fortislifesciences.com/p/A50-201D2>

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: October 24, 2023

Phone: 800.338.9579 • Fax: 866.597.6105 • Web: www.fortislifesciences.com
Orders: orders@fortislifesciences.com • Support: technical@fortislifesciences.com



Detection of hnRNP, Actin, and GAPDH in HeLa Whole Cell Lysate. *Primary Antibodies:* cocktail of mouse anti-hnRNP A500-011A (A500-011A-1), rabbit anti-actin A300-491A (A300-491A-5), and goat anti-GAPDH A303-878A (A303-878A-1) at 1 µg/ml each. *Secondary Antibodies:* cocktail of Dylight® 680-conjugated donkey anti-mouse A90-337D6-2 (A90-337D6-2) (green), Dylight® 800-conjugated donkey anti-rabbit A120-208D8 (A120-208D8-1) (red), and Dylight® 488-conjugated donkey anti-goat A50-201D2 (A50-201D2-4) (blue) at 0.5 µg/ml each. *Acquisition:* Syngene G:Box, 73 seconds (green), 45 seconds (red), and 25 seconds (blue).