Horse IgG-heavy and light chain Antibody

Goat Polyclonal Conjugate DyLight® 488

Antigen Affinity Purified Catalog No. A70-106D2 A70-106D2-6 Lot No.

APPLICATIONS IHC, ICC, F, IF

SPECIES REACTIVITY Horse **AMOUNT** 1 ml

CONCENTRATION 0.5 ma/ml

2 - 8° C / 1 year from date of receipt STORAGE/SHELF LIFE

PHYSICAL STATE Liquid

BUFFER Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide

FLUOROPHORE/PROTEIN 5.9 **ISOTYPE** IqG USA **ORIGIN**

PRODUCTION The antibody to horse IgG was isolated by affinity chromatography using antigen coupled to **PROCEDURES**

agarose beads and conjugated to DyLight® 488.

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of

1.4 equals 1.0 mg of IgG.

By immunoelectrophoresis and ELISA this antibody reacts specifically with horse IgG and with light chains common to other horse immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from

other species.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry 1:50 - 1:500 Immunocytochemistry 1:50 - 1:500

Flow Cytometry 1:50 - 1:200 Immunofluorescence 1:50 - 1:500

APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.

DyLight® 488 is excited at 493 (in PBS) and emits at 518 (in PBS).

DvLight[®] is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.

ADDITIONAL INFO https://www.bethyl.com/product/A70-106D2

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush. PhD | Chief Scientific Officer Date: Iulv 16, 2020

