Granzyme B Recombinant Monoclonal Antibody [BLR022E]





Rabbit Recombinant Monoclonal

Purified Protein ID NP_004122.2

Catalog No. A700-022-T GeneID 3002

Lot No. A700-022-T-3

APPLICATIONS WB, IHC, ICC, IHC-IF, mIF

SPECIES REACTIVITY Human

AMOUNT 10 μ l (5+ tests)

CONCENTRATION 50 μg/ml

STORAGE/SHELF LIFE 2 – 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Borate Buffered Saline (BBS) pH 8.2 with 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG

CLONE # BLR022E

ORIGIN USA

PRODUCTION

The monoclonal antibody was purified from cell culture supernatant.

PROCEDURES

Immunogen was a recombinant protein representing human Granzyme B (residues 19-247,

NP_004122.2).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1000

Immunohistochemistry 1:100 – 1:500. Epitope retrieval with citrate buffer pH 6.0 is

recommended for FFPE tissue sections.

Immunocytochemistry 1:100 - 1:500. Epitope retrieval with citrate buffer pH6.0 is

recommended for FFPE cell sections.

Immunofluorescence 1:100 to 1:500. Epitope retrieval with citrate buffer pH6.0 is

(IHC) recommended for FFPE tissue sections.

Multiplex 1.250

Immunofluorescence

IHC HUMAN CONTROLS Appendix, Breast Carcinoma, Lung Carcinoma, Ovarian Carcinoma, Prostate Carcinoma, Skin

Carcinoma, Spleen, Tonsil, SR cells


ADDITIONAL INFO https://www.bethyl.com/product/A700-022-T


Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: April 12, 2022






Detection of human Granzyme B by western blot.


Samples: Whole cell lysate (15 µg) from HEK293T, HeLa, SR, Jurkat, and Hep-G2 cells prepared using NETN lysis buffer. *Antibody:* Rabbit anti-GranzymeB recombinant monoclonal antibody [BLR022E] (A700-022-T lot 3) used at 1:1000. *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-101P). Chemiluminescence with an exposure time of 10 seconds. Lower Panel: Rabbit anti-COPB2 antibody (A304-523A).

Detection of human Granzyme B by western blot.


Samples: Myc-tagged Granzyme H and B overexpressing cell lysate (15 µg) prepared using NETN lysis buffer. Antibody: Rabbit anti-Granzyme B recombinant monoclonal antibody [BLR022E] (A700-022-T lot 3) used at 1:1000. Secondary: HRP-conjugated goat anti-rabbit IgG (A120-101P). Chemiluminescence with an exposure time of 3 seconds. Lower Panel: Rabbit anti-myc antibody (A191-101).




Detection of human Granzyme B by immunohistochemistry. *Sample:* FFPE section of lymph node. *Antibody:* Rabbit anti-Granzyme B recombinant monoclonal antibody [BLR022E] (A700-022-T lot 3). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P).



Detection of human Granzyme B by immunohistochemistry. *Sample:* FFPE section of lung carcinoma. *Antibody:* Rabbit anti–Granzyme B recombinant monoclonal antibody [BLR022E] (A700–022–T lot 3). *Secondary:* HRP-conjugated goat anti–rabbit IgG (A120–501P).



Detection of human Granzyme B by immunocytochemistry. *Sample:* FFPE section of Granzyme B over-expressing cells (left) and Granzyme H over-expressing cells (right). *Antibody:* Rabbit anti-Granzyme B recombinant monoclonal antibody [BLR022E] (A700-022-T lot 3). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P).



Detection of human Granzyme B (red) by immunohistochemistry. Sample: FFPE section of human lung carcinoma. Antibody: Rabbit anti-Granzyme B recombinant monoclonal antibody [BLR022E] (A700-022-T lot 1) used at 1:250. Secondary: HRP-conjugated goat antirabbit IgG (A120-501P). Substrate: Opal™. Counterstain: DAPI (blue).