## FOXP3 Recombinant Monoclonal Antibody [BLR034F]



Rabbit Recombinant Monoclonal

Purified Protein ID NP\_054728.2

Catalog No. A700-034 GeneID 50943

Lot No. A700-034-2

**APPLICATIONS** WB, IP, IHC, ICC, IHC-IF, mIF

**SPECIES REACTIVITY** Human

**AMOUNT** 100 µl (50+ tests)

CONCENTRATION  $1000 \mu g/ml$ 

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

Borate Buffered Saline (BBS) pH 8.2 with 0.09% Sodium Azide, BSA Free **BUFFER** 

**ISOTYPE** 

CLONE # BLR034F

**ORIGIN USA** 

**PRODUCTION PROCEDURES**  Recombinant antibody was purified from cell culture supernatant.

Immunogen was a peptide representing a region between residue 275 and 325 of human

Forkhead box P3 using the numbering given in entry NP\_054728.2 (Gene ID 50943).

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1000

Immunoprecipitation 20 ul/mg lysate

Immunohistochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH6.0 is

recommended for FFPE tissue sections.

Immunocytochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH6.0 is

recommended for FFPE cell sections.

Immunofluorescence

(IHC)

1:100 to 1:500. Epitope retrieval with citrate buffer pH6.0 is

recommended for FFPE tissue sections.

Multiplex 1:250


Immunofluorescence


**IHC HUMAN CONTROLS** Breast Carcinoma, Lung Carcinoma, Prostate Carcinoma, Renal Cell Carcinoma, Tonsil, MJ Cells

**ADDITIONAL INFO** https://www.fortislife.com/p/A700-034

Use the link above to view SDS, a current list of citations, and other product specific information.


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Michael Spencer, PhD Date: August 5, 2022






Detection of human FOXP3 by western blot. Samples: Whole cell lysate (50 μg) from Jurkat, HEK293T, MJ, MOLT-4, and RPMI-8226 cells prepared using NETN lysis buffer. Antibody: Rabbit anti-FOXP3 recombinant monoclonal antibody [BLR034F] (A700-034 lot 2) used at 1:1000. Secondary: HRP-conjugated goat anti-rabbit IgG (A120-101P). Chemiluminescence with an exposure time of 30 seconds. Lower Panel: Rabbit anti-COPB2 antibody (A304-523A).

Detection of human FOXP3 by western blot of immunoprecipitates. Samples: Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from MJ cells prepared using NETN lysis buffer. Antibodies: Rabbit anti-FOXP3 recombinant monoclonal antibody [BLR034F] (A700-034 lot 2) used for IP at 20 µl/mg lysate. FOXP3 was also immunoprecipitated by a previous lot of this antibody (A700-034 lot 1). For blotting immunoprecipitated FOXP3, A700-034 was used at 1:1000. Chemiluminescence with an exposure time of 30 seconds.



**Detection of human FOXP3 in FFPE tonsil by immunohistochemistry.** *Antibody:* Rabbit anti-FOXP3 recombinant monoclonal antibody [BLR034F] (A700–034 lot 2). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120–501P). *Substrate:* DAB.

**Detection of human FOXP3 in FFPE breast carcinoma by immunohistochemistry.** *Antibody:* Rabbit anti-FOXP3 recombinant monoclonal antibody [BLR034F] (A700-034 lot 2). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P). *Substrate:* DAB.



**Detection of human FOXP3 in FFPE MJ cells by immunocytochemistry.** *Antibody:* Rabbit anti–FOXP3 recombinant monoclonal antibody [BLR034F] (A700–034 lot 2). *Secondary:* HRP–conjugated goat anti–rabbit IgG (A120–501P). *Substrate:* DAB.

Detection of human FOXP3 (red) by immunohistochemistry. Sample: FFPE section of human breast carcinoma. Antibody: Rabbit anti-FOXP3 recombinant monoclonal antibody [BLR034F] (A700-034 lot 1) used at 1:250. Secondary: HRP-conjugated goat antirabbit IgG (A120-501P). Substrate: Opal™. Counterstain: DAPI (blue).