
## VISTA Recombinant Monoclonal Antibody [BLR035F]




Rabbit Recombinant Monoclonal					
Purified			Protein ID	NP_071436.1	
Catalog No. A700-035-T		035-T	GenelD	64115	
Lot No.	A700-0	035-T-2			
APPLICATIONS		WB, IP, IHC, ICC, IHC-IF, Flow Cyt			
SPECIES REACTIVITY		Human			
AMOUNT		10 µl (5+ tests)			
CONCENTRATION		100 μg/ml			
STORAGE/SHELF LIFE		2 - 8°C / 1 year from date of receipt			
PHYSICAL STATE		Liquid			
BUFFER		Borate Buffered Saline (BBS) pH 8.2 with 0.1% BSA and 0.09% Sodium Azide			
ISOTYPE		IgG			
CLONE #		BLR035F			
ORIGIN		USA			
PRODUCTION PROCEDURES		Recombinant antibody was purified from cell culture supernatant.			
		Immunogen was a peptide representing a region between residue 250 and the C-terminus (residue 311) of human V-type immunoglobulin domain-containing suppressor of T-cell activation precursor using the numbering given in entry NP_071436.1 (Gene ID 64115).			
APPLICATIONS		Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.			
		Western Blot	1	:1000	
		Immunoprecipita	ation 2	0 ul/1mg lysate	
		Immunohistoche		:100 – 1:500. Epitope retrieval with citrate buffer pH6.0 is ecommended for FFPE tissue sections.	
		Immunocytoche		:100 to 1:500. Epitope retrieval with citrate buffer pH6.0 is ecommended for FFPE cell sections.	
		Immunofluoresc (IHC)	-	:100 to 1:500. Epitope retrieval with citrate buffer pH6.0 is ecommended for FFPE tissue sections.	
		Flow Cytometry		ixed in 4% formaldehyde and permeabilized with 90% methanol. 2 $\mu l$ er 1 x 10^6 cells.	
IHC HUMAN CONTROLS		Colon Carcinoma, Lung Carcinoma, Placenta, NCI-H226 Cells			
ADDITIONAL INFO		https://www.fortislife.com/p/A700-035-T Use the link above to view SDS, a current list of citations, and other product specific information.			

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Michael Spencer, PhD Date: August 5, 2022

> Phone: 800.338.9579 · Fax: 866.597.6105 · Web: www.fortislife.com Orders: orders@fortislife.com · Support: technical@fortislife.com



Detection of human VISTA by western blot. Samples: Whole cell lysate (50 µg) from HeLa, Jurkat, HEK293T, KG-1, and RPMI-8226 cells prepared using NETN lysis buffer. Antibody: Rabbit anti-VISTA recombinant monoclonal antibody [BLR035F] (A700-035-T lot 2) used at 1:1000. Secondary: HRP-conjugated goat anti-rabbit IgG (A120-101P). Chemiluminescence with an exposure time of 10 seconds. Lower Panel: Rabbit anti-Actin recombinant monoclonal antibody [BLR057F] (A700-057).



Detection of human VISTA by western blot of immunoprecipitates. Samples: Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from Jurkat cells prepared using NETN lysis buffer. Antibodies: Rabbit anti-VISTA recombinant monoclonal antibody [BLR035F] (A700-035-T lot 2) used for IP at 10 µl/mg lysate. VISTA was also immunoprecipitated by a second antibody against a different epitiope of VISTA (BL-435-5G11). For blotting immunoprecipitated VISTA, A700-035-T was used at 1:1000. Chemiluminescence with an exposure time of 30 seconds.


Detection of human VISTA by immunohistochemistry. Sample: FFPE section of lung carcinoma. Antibody: Rabbit anti-VISTA recombinant monoclonal antibody [BLR035F] (A700-035-T lot 2) used at 1:250. Secondary: HRP-

conjugated goat anti-rabbit IgG (A120-501P).

## Detection of human VISTA by immunocytochemistry.


Sample: FFPE section of NCI-H226 cells. Antibody: Rabbit anti-VISTA recombinant monoclonal antibody [BLR035F] (A700-035-T lot 2) used at 1:250. Secondary: HRPconjugated goat anti-rabbit IgG (A120-501P).

For in vitro laboratory use only. Not for any clinical, therapeutic or diagnostic use in humans or animals. Not for human or animal consumption. This product may not be resold or modified for resale without the prior written approval of Bethyl Laboratories, Inc. The information provided in this data sheet is believed to be correct but does not purport to be all-inclusive and is intended to be used as a guide. Bethyl Laboratories, Inc. shall not be liable or responsible in any way for use of either this information or the material supplied. Disposal of hazardous material may be subject to federal, state or local laws or regulations.



## Detection of human VISTA (red) by

immunohistochemistry. Sample: FFPE section of human ovarian carcinoma. Antibody: Rabbit anti-VISTA recombinant monoclonal antibody [BLR035F] (A700-035-T lot 1) used at 1:250. Secondary: HRP-conjugated goat antirabbit IgG (A120-501P). Substrate: Opal<sup>™</sup>. Counterstain: DAPI (blue).



Detection of human VISTA in human peripheral blood mononuclear cells by flow cytometry. *Antibody:* Rabbit anti-VISTA recombinant monoclonal [BLR035F] (A700-035-T lot 1) (right) or isotype control (left). *Secondary:* DyLight<sup>®</sup> 488-conjugated goat anti-rabbit IgG (A120-101D2).