CD33 Recombinant Monoclonal Antibody [BLR061G]

Rabbit Recombinant Monoclonal

Purified Protein ID NP_001763.3

Catalog No. A700-061-T GeneID 945

Lot No. A700-061-T-1

APPLICATIONS WB, IP, IHC

SPECIES REACTIVITY Human

AMOUNT 10 μ l (5+ tests)

CONCENTRATION 100 μg/ml

STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Borate Buffered Saline (BBS) pH 8.2 with 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG

CLONE # BLR061G
ORIGIN USA

PRODUCTION PROCEDURES

Recombinant antibody was purified from cell culture supernatant.

Immunogen was a peptide representing a region between residue 314 and the C-terminus (residue 364) of human Myeloid cell surface antigen CD33 using the numbering given in entry

NP_001763.3 (Gene ID 945).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1000

Immunoprecipitation 20 μ I/1 mg lysate

Immunohistochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH6.0 is

recommended for FFPE sections.


IHC HUMAN CONTROLS Acute Myeloid Leukemia (AML)

ADDITIONAL INFO https://www.bethyl.com/product/A700-061-T


Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019






Detection of human CD33 by immunohistochemistry. *Sample:* FFPE section of Acute Myeloid Leukemia (AML). *Antibody:* Rabbit anti-CD33 recombinant monoclonal antibody [BLR061G] (A700-061-T lot 1) used at 1:200. *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P). *Substrate:* DAB.



Detection of human CD33 by western blot. Samples: Whole cell lysate (50 μg) from HEL 92.1.7, Jurkat, My-La CD4+, HEK293T, HeLa, and THP-1 cells prepared using NETN lysis buffer. Antibody: Rabbit anti-CD33 recombinant monoclonal antibody [BLR061G] (A700-061-T lot 1) used at 1:1000. Secondary: HRP-conjugated goat anti-rabbit IgG (A120-101P). Chemiluminescence with an exposure time of 30 seconds. Lower Panel: Rabbit anti-GAPDH (A300-639A).



Detection of human CD33 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from HEL 92.1.7 cells prepared using NETN lysis buffer. *Antibodies:* Rabbit anti-CD33 recombinant monoclonal antibody [BLR061G] (A700–061–T lot 1) used for IP at 20 μI/mg lysate. CD33 was also immunoprecipitated by rabbit anti-CD33 recombinant monoclonal antibody [BL-302–3E10]. For blotting immunoprecipitated CD33, A700–061–T was used at 1:1000. Chemiluminescence with an exposure time of 10 seconds.