SMARCA2/BRM Recombinant Monoclonal Antibody [BLR302M]



Rabbit Recombinant Monoclonal

Purified RefSeq ID NP_003061.3

Catalog No. A700-302-T Uniprot ID P51531 Lot No. 1 GeneID 6595

APPLICATIONS IHC, ICC, Flow Cyt

SPECIES REACTIVITY Human, Mouse

AMOUNT 10 μ l (5+ tests)

CONCENTRATION 250 μg/ml

STORAGE/SHELF LIFE 2 – 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Phosphate Buffered Saline (PBS) with 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG

CLONE # BLR302M

ORIGIN USA

PRODUCTION PROCEDURES

Recombinant antibody was purified from cell culture supernatant.

Immunogen was a peptide representing a region between residue 100 and 150 of human

SMARCA2/BRM using the numbering given in entry NP_003061.3 (Gene ID 6595).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is

recommended for FFPE tissue sections.

Immunocytochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is

recommended for FFPE cell sections.

Flow Cytometry Fixed in 4% formaldehyde and permeabilized with 90% methanol. 1 µl

per 1 x 10^6 cells.

IHC HUMAN CONTROLS Breast Carcinoma, Ovarian Carcinoma, Prostate Carcinoma, Skin Carcinoma, Tonsil, A-431 Cells,

Cal-27 Cells, KG-1 Cells, VLN3G2 Cells

IHC MOUSE CONTROLS Kidney, Lung, Spleen, Testis, BW5147.3 Cells, J774A1 Cells, NIH 3T3 Cells


ADDITIONAL INFO https://www.fortislife.com/p/A700-302-T

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD


Date: December 26, 2023



Detection of human SMARCA2/BRM by immunohistochemistry. Sample: FFPE section of ovarian carcinoma. Antibody: Rabbit anti-SMARCA/BRM recombinant monoclonal antibody [BLR302M] (A700-302). Secondary: HRP-conjugated goat anti-rabbit IgG (A120-501P).




Detection of human SMARCA2/BRM by immunocytochemistry. *Sample:* FFPE section of VLN3G2 cells. *Antibody:* Rabbit anti-SMARCA/BRM recombinant monoclonal antibody [BLR302M] (A700–302). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120–501P).




Detection of human SMARCA2/BRM (shaded) in HeLa cells by flow cytometry. *Antibody:* Rabbit anti-SMARCA2/BRM recombinant monoclonal antibody [BLR302M] (A700–302) or isotype control (unshaded). *Secondary:* DyLight® 650–conjugated goat anti-rabbit IgG (A120–101D5).



Detection of mouse SMARCA2/BRM by immunohistochemistry. *Sample:* FFPE section of kidney. *Antibody:* Rabbit anti-SMARCA/BRM recombinant monoclonal antibody [BLR302M] (A700–302). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120–501P).



Detection of mouse SMARCA2/BRM by immunocytochemistry. *Sample:* FFPE section of J774A.1 cells. *Antibody:* Rabbit anti-SMARCA/BRM recombinant monoclonal antibody [BLR302M] (A700–302). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120–501P).



Detection of mouse SMARCA2/BRM (shaded) in NIH3T3 cells by flow cytometry. *Antibody:* Rabbit anti-SMARCA2/BRM recombinant monoclonal antibody [BLR302M] (A700–302) or isotype control (unshaded). *Secondary:* DyLight® 650-conjugated goat anti-rabbit IgG (A120–101D5).