

# Human IgG–Fc Fragment Antibody

Goat Polyclonal Conjugate Biotin

Antigen Affinity Purified

Catalog No. A80–104B

Lot No. 19

**APPLICATIONS** WB, IHC, ICC, ELISA

**SPECIES REACTIVITY** Human

**AMOUNT** 1 ml

**CONCENTRATION** 1 mg/ml

**STORAGE/SHELF LIFE** 2 – 8°C / 1 year from date of receipt

**PHYSICAL STATE** Liquid

**BUFFER** Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide

**ISOTYPE** IgG

**ORIGIN** USA

**PRODUCTION PROCEDURES** Antiserum was solid phase adsorbed to ensure class specificity. The antibody was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to biotin.

Antibody concentration was determined by extinction coefficient prior to conjugation: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

By immunoelectrophoresis and ELISA this antibody reacts specifically with human IgG. Cross reactivity with IgA, IgM and light chains is less than 1%. This antibody may cross react with IgG from other species.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:20,000 – 1:400,000

Immunohistochemistry 1:500 – 1:5,000

Immunocytochemistry 1:200 – 1:1,000

ELISA 1:20,000 – 1:400,000

**APPLICATION NOTES** Not all listed applications have been specifically tested by our laboratory.

**ADDITIONAL INFO** [https://www.fortislife.com/p/A80–104B](https://www.fortislife.com/p/A80-104B)

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.  
Michael Spencer, PhD Date: June 27, 2023