Mouse IgG Heavy and Light Chain Antibody



Goat Polyclonal Conjugate DyLight® 650

Antigen Affinity Purified Catalog No. A90-116D5

Lot No. 11

APPLICATIONS IHC, ICC, Flow Cyt, IF

SPECIES REACTIVITY Mouse
AMOUNT 1 ml

CONCENTRATION 0.5 mg/ml

STORAGE/SHELF LIFE 2 – 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide

FLUOROPHORE/PROTEIN 6.2
ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

The antibody was isolated by affinity chromatography using antigen coupled to agarose

beads and conjugated to DyLight® 650.

Immunoglobulin concentration was determined using Beer's Law where 1 mg/mL lgG has an

A280 of 1.4.

By immunoelectrophoresis and ELISA this antibody reacts specifically with mouse IgG and with light chains common to other mouse immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from

other species.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry1:50 - 1:500Immunocytochemistry1:50 - 1:500Flow Cytometry1:50 - 1:200Immunofluorescence1:50 - 1:500

APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.

DyLight® 650 is excited at 652 (in PBS) and emits at 672 (in PBS). DyLight® 650 replaces DyLight®

649.

DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.

ADDITIONAL INFO https://www.fortislife.com/p/A90-116D5


Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: January 19, 2024

Phone: 800.338.9579 • Fax: 866.597.6105 • Web: www.fortislife.com Orders: orders@fortislife.com • Support: technical@fortislife.com



Detection of human HLA-DR (shaded) in Daudi cells by flow cytometry. *Antibody:* Mouse anti-HLA-DR monoclonal antibody [LN3] (A500-022A) or isotype control (unshaded). *Secondary:* DyLight® 650-conjugated goat anti-mouse IgG (A90-116D5).