

Mouse IgG Heavy and Light Chain Antibody

Donkey Polyclonal Conjugate DyLight® 594

Antigen Affinity Purified

Catalog No. A90-137D4

Lot No. A90-137D4-7

APPLICATIONS	IHC, ICC, F, IF
SPECIES REACTIVITY	Mouse
AMOUNT	1 ml
CONCENTRATION	0.5 mg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide
FLUOROPHORE/PROTEIN	6.9
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	The antibody was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight® 594.

Antibody concentration was determined by extinction coefficient prior to conjugation: absorbance at 280 nm of 1.4 equals 1.0 mg of gG.

By immunoelectrophoresis and ELISA this antibody reacts specifically with mouse IgG and with light chains common to other mouse immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry	1:40 – 1:400
Immunocytochemistry	1:40 – 1:400
Flow Cytometry	1:50 – 1:200
Immunofluorescence	1:40 – 1:400

APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.


DyLight® 594 is excited at 593 (in PBS) and emits at 618 (in PBS).

DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.

ADDITIONAL INFO <https://www.bethyl.com/product/A90-137D4>

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: March 18, 2022



Detection of human p53 by immunofluorescence. *Sample:* FFPE serial sections of human stomach carcinoma. *Primary Antibody:* mouse anti-p53 (Clone DO-1) used at a dilution of 1:100. *Secondary Antibody:* Red-fluorescent donkey anti-mouse IgG-heavy and light chain Antibody DyLight® 594 Conjugated (A90-137D4 Lot 3) used at a dilution of 1:100 (5µg/ml). *Counterstain:* DAPI (blue)