Mouse IgG-F(ab')2 Fragment Antibody



F(ab')2 Goat Polyclonal Conjugate

Antigen Affinity Purified A90-141F Catalog No. Lot No. A90-141F-13

IHC, ICC, F, IF **APPLICATIONS**

SPECIES REACTIVITY Mouse **AMOUNT** 1 ml

CONCENTRATION 0.5 mg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide

FITC

FLUOROPHORE/PROTEIN 4.7 **ISOTYPE** IqG **ORIGIN** USA

PRODUCTION PROCEDURES

Antiserum was solid phase adsorbed to ensure class specificity. The antibody to mouse IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Fragments were conjugated to fluorescein isothiocyanate (FITC).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

By immunoelectrophoresis and ELISA this antibody reacts specifically with Fab fragment of mouse IgG and with light chains common to other mouse immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react

with Fab fragments from other species.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

1:50 - 1:500

experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry 1:50 - 1:500 Immunocytochemistry 1:50 - 1:500 Flow Cytometry 1:50 - 1:200Immunofluorescence

APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.

ADDITIONAL INFO https://www.bethyl.com/product/A90-141F

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Michael Spencer, PhD Date: November 9, 2021