

Mouse IgG-heavy and light chain cross-adsorbed Antibody

Rabbit Polyclonal Conjugate DyLight® 594
Antigen Affinity Purified
Catalog No. A90-317D4
Lot No. A90-317D4-7



APPLICATIONS	IHC, ICC, F, IF
SPECIES REACTIVITY	Mouse. Minimum reactivity to human and rat
ISOTYPE	IgG
AMOUNT	1 ml at 0.5 mg/ml
STORAGE/SHELF LIFE	2 - 8° C / 1 year from date of receipt
PHYSICAL STATE	Liquid
FLUOROPHORE/PROTEIN	4.2
BUFFER	Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide
ORIGIN	USA
PRODUCTION PROCEDURES	Antiserum was cross adsorbed using human and rat immunosorbents to remove cross reactive antibodies. The antibody to mouse IgG was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight® 594.

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

By immunoelectrophoresis and ELISA this antibody reacts specifically with mouse IgG and with light chains common to other mouse immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. Less than 1% cross reactivity to human and rat IgG was detected. This antibody may cross react with IgG from other species.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry	1:50 – 1:500
Immunocytochemistry	1:50 – 1:500
Flow Cytometry	1:50 – 1:200
Immunofluorescence	1:50 – 1:500

APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.

DyLight® 594 is excited at 593 (in PBS) and emits at 618 (in PBS).


DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.

ADDITIONAL INFO <https://www.bethyl.com/product/A90-317D4>
Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Eric McIntush, PhD | Chief Scientific Officer

Date: December 3, 2018



Detection of human p53 by immunofluorescence. *Sample:* FFPE serial sections of human stomach carcinoma. *Primary Antibody:* mouse anti-p53 (Clone DO-1) used at a dilution of 1:100. *Secondary Antibody:* Red-fluorescent Rabbit anti-mouse IgG-heavy and light chain cross-adsorbed Antibody DyLight® 594 Conjugated (A90-317D4 Lot 6) used at a dilution of 1:100 (5µg/ml). *Counterstain:* DAPI (blue)