Mouse IgG-heavy and light chain cross-adsorbed Antibody BETHYL





DyLight® 594 **Donkey Polyclonal** Conjugate

Antigen Affinity Purified A90-337D4 Catalog No. A90-337D4-14 Lot No.

APPLICATIONS IHC, ICC, F, IF

SPECIES REACTIVITY . Minimum reactivity to bovine, chicken, goat, human, rabbit, rat and sheep

AMOUNT

CONCENTRATION 0.5 mg/ml

2 - 8° C / 1 year from date of receipt STORAGE/SHELF LIFE

PHYSICAL STATE Liquid

Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide BUFFER

FLUOROPHORE/PROTEIN 5.7 **ISOTYPE** IqG **ORIGIN** USA

PRODUCTION PROCEDURES

Antiserum was cross adsorbed using bovine, chicken, goat, human, rabbit, rat and sheep immunosorbents to remove cross reactive antibodies. The antibody to mouse IgG was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight® 594.

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

By immunoelectrophoresis and ELISA this antibody reacts specifically with mouse IgG and with light chains common to other mouse immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. Less than 2% cross reactivity to bovine, chicken, goat, human, rabbit, rat and sheep IgG was detected. This antibody may cross react with IaG from other species.

APPLICATIONS

Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry 1:50 - 1:500 Immunocytochemistry 1:50 - 1:500Flow Cytometry 1:50 - 1:200Immunofluorescence 1:50 - 1:500

APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.

DyLight® 594 is excited at 593 (in PBS) and emits at 618 (in PBS).


DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.

ADDITIONAL INFO https://www.bethyl.com/product/A90-337D4

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Michael Spencer, PhD Date: September 1, 2021

> Phone: 800.338.9579 • Fax: 866.597.6105 • Web: www.bethyl.com Orders: orders@fortislife.com · Support: technical@fortislife.com



Detection of human p53 by immunofluorescence. Samples: FFPE serial sections of human stomach carcinoma. Primary Antibody: mouse anti-p53 (Clone DO-1) used at a dilution of 1:100. Secondary Antibody: Red-fluorescent donkey anti-mouse IgG-heavy and light

chain cross-adsorbed Antibody DyLight® 594 Conjugated A90-337D4 Lot 4 (left) and A90-337D4 Lot 5 (right) used at a dilution of 1:100 (5µg/ml). *Counterstain:* DAPI (blue)