PARP1 IHC Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_001609.1

Catalog No. IHC-00279-T GeneID 142

Lot No. IHC-00279-T-1

APPLICATIONS IHC, IHC–IF

SPECIES REACTIVITY Human

AMOUNT 10 μl

CONCENTRATION 250 μg/ml

STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to PARP1 immobilized on solid support.

The epitope recognized by IHC-00279-T maps to a region between residue 50 and 100 of human poly (ADP-ribose) polymerase family, member 1 using the numbering given in entry

NP_001609.1 (GeneID 142).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry 1:100 – 1:500
Immunofluorescence 1:50 – 1:500

(IHC)

APPLICATION NOTES Epitope exposure is recommended.

Epitope exposure with citrate buffer will enhance staining.

Likely to work with frozen sections.

In some cases, the antibody may be diluted further than indicated.


IHC HUMAN CONTROLS Breast Carcinoma, Ovarian Carcinoma

ADDITIONAL INFO https://www.bethyl.com/product/IHC-00279-T


Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019






Detection of human PARP1 by immunohistochemistry. Sample: FFPE section of human breast carcinoma. Antibody: Affinity purified rabbit anti-PARP1 (Cat. No. IHC-00279-T) used at a dilution of 1:250. Detection: DAB



Detection of human PARP1 by immunohistochemistry. Sample: FFPE section of human ovarian carcinoma. Antibody: Affinity purified rabbit anti-PARP1 (Cat. No. IHC-00279-T) used at a dilution of 1:250. Detection: DAB



Detection of human PARP1 by immunohistochemistry. Sample: FFPE section of human breast carcinoma. Antibody: Affinity purified rabbit anti-PARP1 (Cat. No. IHC-00279-T) used at a dilution of 1:100. Detection: Redfluorescent goat anti-rabbit IgG highly cross-adsorbed Antibody used at a dilution of 1:100.